

CocciAid[®] for Calves (amprolium)

9.6% Oral Solution Coccidiostat

INDICATIONS FOR USE: An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

DOSAGE AND ADMINISTRATION:

5-Day Treatment Protocol

Daily Dosage: 10 mg amprolium/kg
(10 mg per 2.2 lb body weight)

21-Day Prevention Protocol

Daily Dosage: 5 mg amprolium/kg
(5 mg per 2.2 lb body weight)

USE DIRECTIONS: There are **96 mg** of amprolium in every **1 mL** of CocciAid for Calves 9.6% Solution.

1 fl oz = 29.57 mL

1 fl oz = 2 measuring tablespoons

8 fl oz = 1/2 pint (236.56 mL)

16 fl oz = 1 pint (473.12 mL)

IN DRINKING WATER

Mix CocciAid for Calves daily with fresh drinking water.

5-DAY TREATMENT: Add CocciAid for Calves 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CocciAid for Calves 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

AS A DRENCH

CocciAid for Calves drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add 3 fl oz CocciAid for Calves 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of CocciAid for Calves 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

Use as the sole source of amprolium.

USER SAFETY WARNINGS: NOT FOR HUMAN USE.

Keep this and all drugs out of the reach of children.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Aurora Pharmaceutical at 1-888-215-1256 or www.aurorapharmaceutical.com. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimalae.

PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION. For irritations, flush with plenty of water; get medical attention.

STORAGE: Store between 5°-25°C (41°- 7°F) with brief excursions to 40°C.

Benzoic acid 0.1% added as preservative.

Restricted Drug (California) - Use Only as Directed

Approved by FDA under ANADA # 200-780

CocciAid for Calves is a registered trademark of Aurora Pharmaceutical, Inc. ©2025

REORDER NO: 24004

Manufactured by:

Aurora Pharmaceutical, Inc.
Northfield, MN 55057



WITHDRAWAL PERIODS AND RESIDUE WARNINGS:

Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.